



Listing of Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application.

1. (Previously presented) A compound of the general formula (I)



in which B is a radical of the general formula (II)



in which

n is an integer from 2 to 8;

m is 0, 1 or 2;

x is an integer from 0 to 8;

y is an integer from 1 to 4;

z is an integer from 0 to 5;

$R_3$  is an alkyl radical having 1 to 3 C atoms, which may be substituted by one or more hydroxyl groups;  
and in which A is:



in which

$p \geq 0$ ;

$q \geq 0$ ;

$12 \leq p + q \leq 30$ ; and

with the proviso that when  $p + q$  is 12,  $q$  is not 4 and when  $p + q = 14, 16, 18$  or  $20$ ,  $q$  is not 8; and wherein, in A, the double bond is at a distance from O which does not appear in a naturally-occurring corresponding radical.

2. (Original) A compound as claimed in claim 1, in which the following applies to B:

$m = 1$ .

3. (Original) A compound as claimed in claim 2, in which the following applies to B:

$m = 1$ ;

$x = 1$  to  $3$ ;

$z = 0$ .

4. (Original) A compound as claimed in claim 3, in which the following applies to B:

$m = 1$ ;

$x = 1$ ;

$z = 0$ .

5. (Original) A compound as claimed in claim 1, in which the following applies to B:

$m = 1$ ;

U.S. Application No. 09/762,006  
Reply to Office Action dated June 15, 2004  
Response dated December 15, 2004

x = 0;  
y = 1;  
z = 1 to 5.

6. (Original) A compound as claimed in claim 5, in which the following applies to B:

m = 1;  
x = 0;  
y = 1;  
z = 1 to 3.

7. (Original) A compound as claimed in claim 1, in which the following applies to B:

m = 1;  
x = 0;  
y = 2 to 4;  
z = 1.

8. (Original) A compound as claimed in claim 1, in which the following applies to B:

m = 0;  
x = 0;  
y = 1;  
z = 1 to 5.

9. (Original) A compound as claimed in claim 1, in which the following applies to B:

m = 0;  
x = 0;  
y = 2 to 4;  
z = 1.

10. (Previously presented) A compound as claimed in claim 1, in which the following applies to B:

$R_3 = CH_3.$

11. (Previously presented) A compound as claimed in claim 1, in which the following applies to B:

$R_3 = 1,2\text{-dihydroxypropyl}.$

12. (Previously presented) A compound as claimed in claim 1, in which the following applies to B:

$n = 2 \text{ to } 6.$

13. (Previously presented) A compound as claimed in claim 1, in which the following applies to B:

$n = 3.$

14. Canceled.

15. (Previously presented) A compound as claimed in claim 1, in which A has 16 to 23 carbon atoms.

- 16-32. Canceled.

33. (Previously presented) A pharmaceutical composition, which comprises an active ingredient as claimed in claim 1, where appropriate together with pharmaceutically acceptable diluents, excipients, carriers and fillers.


34-42. Canceled.

43. (Previously presented) A compound according to claim 1, wherein p is 9, q is 8, z is 0, x is 1, m is 1, n is 4 and R<sub>3</sub> is methyl.

44. (Previously presented) A compound of the general formula (I)

(I) A - PO<sub>3</sub> - B

in which B is a radical of the general formula (II)



in which

n is an integer from 4 to 8;

m is 1

x is 1;

z is 0;

R<sub>3</sub> is an alkyl radical having 1 C atoms, which is not substituted by a hydroxyl group;  
and in which A is:



in which

$p \geq 0$ ;

$q \geq 0$ ;


$12 \leq p + q \leq 30$  and

where  $q \neq 8$  for  $p + q = 14, 16, 18$  or  $20$  and wherein, in A, the double bond is at a distance from O which does not appear in a naturally-occurring corresponding radical.

45. (Previously presented) A compound of the general formula (I)

(I)  $A - PO_3 - B$

in which B is a radical of the general formula (II)



in which

n is an integer from 2 to 8

m is 0, 1 or 2;

x is an integer from 0 to 8;

y is an integer from 1 to 4;

z is an integer from 0 to 5;

U.S. Application No. 09/762,006  
Reply to Office Action dated June 15, 2004  
Response dated December 15, 2004

$R_3$  is an alkyl radical having 1 to 3 C atoms, which may be substituted by one or more hydroxyl groups;  
and in which A is:



in which

$$p \geq 0;$$

$$q \geq 0;$$

$$12 \leq p + q \leq 30 \text{ and}$$

with the proviso that  $p + q$  is not 12, 13, 14 or 15 and when  $p + q = 16, 18$  or  $20$ ,  $q$  is not 8, and wherein, in A, the double bond is at a distance from O which does not appear in a naturally-occurring corresponding radical.